



Salmon Oil & Cod Liver Oil

Benefits of Salmon Oil and Cod Liver Oil

Salmon Oil

Children need fish oil to maintain good health. Salmon oil is a natural source of Omega-3, especially rich in EPA & DHA. These nutrients in Salmon oil are very important to contribute healthy heart functions, joint flexibility as well as supporting children brain development, nerve, and visual functions. ^[1]

Cod Liver Oil

The pure cod liver oil derived from the cod fish which is a natural good source of Vitamin A and Vitamin D. ^[2]

Vitamin A is an anti-infective vitamin, beneficial for vision health, protecting our mucous membranes, especially the tissue linings in digestive, respiratory and urinary tracts. Night blindness, dry eyes, eye infections and skin problems are just some of the conditions associated with Vitamin A deficiency. ^[3]

Cod liver oil is also rich in Vitamin D, an important vitamin required for the absorption and utilization of Calcium. It is especially important for the normal growth and development of bones and teeth in children. A severe deficiency of Vitamin D can cause rickets in children. ^[4]

Health Benefits of Fish Oil

- Maintains good health of the retina, cornea and eye-sight, prevents night blindness ^[5]
- Helps the growth of baby's brain, enhance IQ & EQ ^[6]
- Enhances the ability of memory and learning ^[6-7]
- Improvement of Dyslexia ^[8]
- Enhances ability of concentration ^[7]
- Strengthens the functions of trachea and lungs ^[9]
- Maintains a good digestive system and good appetite ^[10]
- Protects the liver, against the pathogen of allergic rhinitis, hyperactive airway and skin allergy ^[11]
- Maintains the health of the skin, hair, nail and the joints ^[12]
- Facilitates the absorption of calcium and phosphorus which are essential in maintaining good health of the bones, teeth, heart and brain ^[12]

Recommended daily dose:

Adults: 2-3 softgel capsules daily

Above 12 years: 1-2 softgel capsules daily

3-12 years: 1 softgel capsule daily or as recommended by physicians.

References:

1. Fats and fatty acids in human nutrition. Report of an expert consultation. Geneva. 10 - 14 November 2008. FAO (Food and Agriculture Organization of the United Nations). Food Nutr Pap. 2010;91:1-166.
2. Green J. Studies on the analysis of vitamins D. 5. The chemical determination of vitamin D in fish-liver oils and other materials containing vitamin A. Biochem J. 1951 Jul;49(2):243-6.
3. Evan Mayo-Wilson, Aamer Imdad, Kurt Herzer, Mohammad Yawar Yakoob, Zulfiqar A Bhutta. Vitamin A supplements for preventing mortality, illness, and blindness in children aged under 5: systematic review and meta-analysis. BMJ. 2011; 343:1-19.
4. Thacher TD, Clarke BL. Vitamin D Insufficiency. Mayo Clin Proc. 2011;86(1):50-60.
5. Querques G, Forte R and Souied EH. Retina and Omega-3. J Nutr Metab. 2011; 2011:748361.
6. Mijna Hadders-Algra. Effect of Long-Chain Polyunsaturated Fatty Acid Supplementation on Neurodevelopmental Outcome in Full-Term Infants. Nutrients 2010 Aug;2(8):790-804.
7. Sinn N, Milte C and Howe PR. Oiling the Brain: A Review of Randomized Controlled Trials of Omega-3 Fatty Acids in Psychopathology across the Lifespan. Nutrients 2010 Feb;2(2):128-170.
8. Richardson AJ, Ross MA. Fatty acid metabolism in neurodevelopmental disorder: a new perspective on associations between attention-deficit/hyperactivity disorder, dyslexia, dyspraxia and the autistic spectrum. Prostaglandin Leukot. Essent. Fatty Acids 2000 Jul-Aug; 63(1-2): 1-9.
9. Tabak C, Wijga AH, de Meer G, Janssen NA, Brunekreef B, Smit HA. Diet and asthma in Dutch school children (ISAAC-2). Thorax. 2006 Dec; 61(12):1048-53.
10. Cencic A and Chingwaru W. The Role of Functional Foods, Nutraceuticals, and Food Supplements in Intestinal Health. Nutrients. 2010 Jun;2(6):611-25.
11. Shapiro H, Tehilla M, Attal-Singer J, Bruck R, Luzzatti R, Singer P. The therapeutic potential of long-chain omega-3 fatty acids in nonalcoholic fatty liver disease. Clin Nutr. 2011 Feb;30(1):6-19.
12. Wann AKT, Mistry J, Knight MM et al. Eicosapentaenoic acid and docosahexaenoic acid reduce interleukin-1b-mediated cartilage degradation. Arthritis Research & Therapy 2010;12(6):R207.